

COMPLETENESS RESULTS FOR MANY-VALUED ŁUKASIEWICZ MODAL SYSTEMS AND RELATIONAL SEMANTICS

GEORGES HANSOUL AND BRUNO TEHEUX

ABSTRACT. The paper is dedicated to the problem of adding a modality to the Łukasiewicz many-valued logics in the purpose of obtaining completeness results for Kripke semantics. We define a class of modal many-valued logics and their corresponding Kripke models and modal many-valued algebras. Completeness results are considered through the construction of a canonical model. Completeness is obtained for modal finitely-valued logics but also for a modal many-valued system with an infinitary deduction rule. We introduce two classes of frames for the finitely-valued logics and show that they define two distinct classes of Kripke-complete logics.

1. INTRODUCTION

Modal logics and many-valued logics were both historically introduced in order to free oneself from the rigidity of propositional logic. With many-valued logics, the logician can choose the truth values of the propositions in a set with more than two elements. With modal logics, the logician introduce a new connector whose aim is, for instance, to model the possibility. Even if these two approaches of the science of reasoning are not born independently (see chapter 21 of [10]), many-valued logics and modal logics received distinguished interests since their birth.

On the one hand, mathematicians tackled many-valued logics (as defined by J. ŁUKASIEWICZ in [14]; see [15] for an English translation and [5] for a monograph on the subject) through their algebraic form: the class of MV-algebras that was introduced by C.C. CHANG in 1958 (see [3] and [4]) in order to obtain an algebraic proof of the completeness result for the infinite-valued ŁUKASIEWICZ logic.

On the other hand, modal logics were also studied through their algebraic disguises, which are the Boolean algebras with operators (introduced in [11] and [12]). But the success of modal logics among the communities of mathematicians, computer scientists and philosophers is a consequence of the relational semantics introduced in the sixties by S. KRIPKE (see [13]). With KRIPKE semantic, also called possible worlds semantic, a formula is possible in a world w if it is true in a world accessible from w . From then on, in their approach of modal logic, mathematicians have been focusing their attention on the connection between the algebraic and the relational semantics. This approach allowed a great simplification of the proof of the completeness of normal modal logics through the construction of the so-called canonical model (see [1] for example).

2000 *Mathematics Subject Classification.* 03B45, 03B50.

Key words and phrases. modal logic, many-valued logic, Kripke semantic, relational semantic, canonical model, MV-algebras.

Since the definition of a KRIPKE model can easily be extended to a many-valued realm, the problem of merging modal and many-valued logic has already been considered by some mathematicians in the last few years (see [8] and [7] or [16] for example). But the algebraic approach and its connection with the relational semantics has never been considered.

In this paper, we introduce some modal many-valued (in a ŁUKASIEWICZ meaning) logics and their corresponding algebras and tackle the problem of the completeness with respect to many-valued KRIPKE models through the construction of a canonical model.

The next section of this paper is dedicated to the introduction of the many-valued KRIPKE models. In the third section, we define the modal many-valued logics. The axiomatization of these logics is very natural since it appears clearly that they admit the class of many-valued KRIPKE models as a sound semantic.

Modal many-valued algebras are introduced in the fourth section as a step toward the construction of the canonical model of a modal many-valued logic, which is the object of the next section. Here, the reader can find one of the main result of this paper: the natural definition of the valuation on the canonical model extends to formulas (Proposition 5.5).

Completeness results with respect to the KRIPKE models of section 2 are considered in section 6. The results are obtained for the finitely-valued logics but also for an infinitary modal many-valued system (i.e. a formal system that admits an infinitary deduction rule). The construction of the canonical model allows us to simplify the axiomatization of the modal finitely-valued logics. We also introduce the class of $n+1$ -frames. Roughly speaking, an $n+1$ -frame is a first-order structure obtained from a frame by deciding to ban some valuations on the frame. We then illustrate the dissimilarity between KRIPKE-completeness and $n+1$ -KRIPKE-completeness.

We conclude the paper by suggesting some tracks for possible projects.

2. A RELATIONAL SEMANTIC FOR $[0, 1]$ -VALUED MODAL LOGICS

The modal many-valued systems that we develop in the sequel of the paper admit a sound (and for some of them complete) relational semantic. Since this semantic is the main strength of these systems, we have decided to first introduce it.

Let us denote by **Prop** an infinite set of propositional variables and by $\mathcal{F}$ the set of formulas defined inductively by the following rules:

- (1) **Prop** $\subseteq \mathcal{F}$;
- (2) if ϕ and ψ are in $\mathcal{F}$ then $\neg\phi$, $\phi \oplus \psi$ and $\Box\phi$ are in $\mathcal{F}$.

The intended meaning of $\phi \oplus \psi$ and $\neg\psi$ is clear (these formulas have their usual ŁUKASIEWICZ meaning) and $\Box\phi$ can be read, for example, as “ ϕ necessary holds”.

Obviously, as usual, we make use of the following abbreviations: the formula $\phi \rightarrow \psi$ stands for $\psi \oplus \neg\phi$, the formula $\psi \odot \phi$ for $\neg(\neg\psi \oplus \neg\phi)$, the formula $\phi \vee \psi$ for $(\phi \rightarrow \psi) \rightarrow \psi$, the formula $\phi \wedge \psi$ for $\neg(\neg\phi \vee \neg\psi)$ and $\Diamond\phi$ for $\neg\Box\neg\phi$.

The definition of a KRIPKE model can easily be extended to a $[0, 1]$ -valued realm. In the following, we consider the real unit interval $[0, 1]$ endowed with the ŁUKASIEWICZ implication and negation: $x \rightarrow y = \min(1, 1 - x + y)$ and $\neg x = 1 - x$. If n is a positive integer, we denote by $\mathbb{L}_n$ the subset $\{0, \frac{1}{n}, \dots, \frac{n-1}{n}, 1\}$ of $[0, 1]$. Note that $\mathbb{L}_n$ is closed for $\rightarrow$ and $\neg$.

Definition 2.1. A *many-valued Kripke model* (or simply a *many-valued model*) $\mathcal{M} = \langle W, R, \text{Val} \rangle$ is given by a non empty set W , an accessibility relation $R \subseteq W \times W$ and a map $\text{Val} : \mathbf{Prop} \times W \rightarrow [0, 1]$. If n is a positive integer such that $\text{Val}(\mathbf{Prop}, W) \subseteq \mathbb{L}_n$, then $\mathcal{M}$ is called an *$n + 1$ -valued Kripke model*.

A *frame* $\mathfrak{F} = \langle W, R \rangle$ is given by an non empty set W and an accessibility relation R on W . A model $\mathcal{M} = \langle W', R', \text{Val} \rangle$ is *based on the frame* $\mathfrak{F} = \langle W, R \rangle$ if $W = W'$ and $R = R'$.

If $\mathcal{M} = \langle W, R, \text{Val} \rangle$ is a many-valued Kripke model, we extend inductively the map Val to formulas of by the following rules:

- $\text{Val}(\phi \oplus \psi, w) = \text{Val}(\phi, w) \oplus \text{Val}(\psi, w)$ and $\text{Val}(\neg \phi, w) = \neg \text{Val}(\phi, w)$,
- $\text{Val}(\Box \phi, w) = \bigwedge \{ \text{Val}(\phi, w') \mid (w, w') \in R \}$,

for any formulas ϕ and ψ of and any world w of W (where $\bigwedge$ means that we consider the infimum in $[0, 1]$).

We write $\mathcal{M}, w \models \phi$ or simply $w \models \phi$ (and say that w *satisfies* ϕ) whenever $\text{Val}(\phi, w) = 1$, and $\mathcal{M} \models \phi$ whenever $w \models \phi$ for any w in W . In that case, we say ϕ is *true* in $\mathcal{M}$. Formulas that are true in any model $\mathcal{M}$ are called *tautologies*. If Γ is a set of formulas that are true in a model $\mathcal{M}$, then $\mathcal{M}$ is a *model of* Γ . If $\mathfrak{F}$ is a frame and ϕ is a formula that is true in any model based on $\mathfrak{F}$, we say that ϕ is *valid* in $\mathfrak{F}$ and write $\mathfrak{F} \models \phi$.

Note that the 2-valued Kripke models coincide with the Kripke models of normal modal logics (where the operation $\oplus$ matches up with the supremum $\vee$).

Proposition 2.2. *If τ is an increasing unary term of the language $\mathcal{L}_{MV} = \{\rightarrow, \neg\}$, then the formulas*

$$\Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q), \quad \Box(p \wedge q) \leftrightarrow \Box p \wedge \Box q, \quad \Box \tau(p) \leftrightarrow \tau(\Box p)$$

are tautologies.

The idea of using these models as a semantic for modal many-valued logics is not new. See [16] for example.

3. ŁUKASIEWICZ MODAL MANY-VALUED LOGICS

The purpose of this section is to introduce a family of *modal many-valued logics* and their corresponding algebras in order to tackle completeness results through the construction of a *canonical model*. We refer to [5] for an introduction to ŁUKASIEWICZ logic and to [1] and [2] for an introduction to modal logic.

Definition 3.1. A *modal many-valued logic* is a set $\mathbf{L}$ of formulas of that is closed under *modus ponens*, substitution, the necessitation rule (RN) (from ϕ infer $\Box \phi$) and that contains

- an axiomatic base of ŁUKASIEWICZ many-valued logic ($p \rightarrow (q \rightarrow p)$, $(p \rightarrow q) \rightarrow ((q \rightarrow r) \rightarrow (p \rightarrow r))$, $((p \rightarrow q) \rightarrow q) \rightarrow ((q \rightarrow p) \rightarrow p)$, $(\neg p \rightarrow \neg q) \rightarrow (q \rightarrow p)$ for example);
- the axiom (K) of modal logic: $\Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$,
- the formulas $\Box(p \oplus p) \leftrightarrow \Box p \oplus \Box p$ and $\Box(p \odot p) \leftrightarrow \Box p \odot \Box p$,
- the formula $\Box(p \oplus p^m) \leftrightarrow (\Box p \oplus (\Box p)^m)$ for every positive integer m .

As usual, we write $\vdash_{\mathbf{L}} \phi$ and say that ϕ is a *theorem of $\mathbf{L}$* whenever $\phi \in \mathbf{L}$ and denote by $\mathbf{K}$ the smallest modal many-valued logic. If in addition $\mathbf{L}$ contains an axiomatic base of the $n + 1$ -valued ŁUKASIEWICZ logic, we say that $\mathbf{L}$ is a *modal $n + 1$ -valued logic* and we denote by $\mathbf{K}_n$ the smallest of these logics.

Note that, according to Proposition 2.2, the proposed axioms are tautologies for the many-valued KRIPKE models, so that the latter form a sound semantic for the modal many-valued logics.

Let us also remark that, as it will appear clearly in the sequel of the paper (in Proposition 5.5), we only use the last family of axioms as a kind of conservative law for $\Box$ with respect to infinitely great elements. Moreover Proposition 6.3 gives an axiomatization of the finitely-valued logics without this family of axioms (and this explains why we have added the axiom $\Box(p \oplus p) \leftrightarrow \Box p \oplus \Box p$ even if it is equivalent to $\Box(p \oplus p^m) \leftrightarrow (\Box p \oplus (\Box p)^m)$ with $m = 1$).

We can easily gather the following theorems and admissible rules of $\mathbf{K}$.

Proposition 3.2. *The following formulas are theorems of $\mathbf{K}$:*

$$\begin{aligned} \Box(p \rightarrow q) &\rightarrow (\Diamond p \rightarrow \Diamond q), & \Diamond(p \oplus q) &\rightarrow (\Diamond p \oplus \Diamond q), \\ (\Box p \wedge \Diamond q) &\rightarrow \Diamond(p \wedge q), & \Box(p \wedge q) &\rightarrow (\Box p \wedge \Box q) \\ (\Box p \odot \Box q) &\rightarrow \Box(p \odot q). \end{aligned}$$

Moreover, the logic $\mathbf{K}$ is closed under the following deduction rules:

$$\begin{aligned} (1) & \frac{\phi \leftrightarrow \psi}{\Box \phi \leftrightarrow \Box \psi}, \\ (2) & \frac{\phi_1 \odot \cdots \odot \phi_n \rightarrow \psi}{\Box \phi_1 \odot \cdots \odot \Box \phi_n \rightarrow \Box \psi}. \end{aligned}$$

Proof. The proofs are simple adaptations of the two-valued proofs. $\square$

Note that at this point of our development, we can not decide if the formula $\Box p \wedge \Box q \rightarrow \Box(p \wedge q)$ is in $\mathbf{K}$ or in $\mathbf{K}_n$ (for $n \geq 2$). We shall conclude latter, thanks to a completeness result, that it is a theorem of $\mathbf{K}_n$ for any n . On the opposite, the formula $\Box(p \odot q) \rightarrow \Box p \odot \Box q$ is not a theorem of $\mathbf{K}_n$ for any $n \geq 2$ since it is not a $\mathbf{K}_n$ -tautology.

Definition 3.3. If $\Gamma \cup \{\phi\}$ is a set of formulas and if $\mathbf{L}$ is a many-valued modal logic, we say that ϕ is *deducible from Γ in $\mathbf{L}$* and write $\Gamma \vdash_{\mathbf{L}} \phi$ (or simply $\Gamma \vdash \phi$ when $\mathbf{L} = \mathbf{K}$ or $\mathbf{L} = \mathbf{K}_n$ following the context) if ϕ is in any extension of $\Gamma \cup \mathbf{L}$ that is closed under substitution and *modus ponens*.

Note that we can state the following adaptation of the deduction theorem.

Lemma 3.4. *If $\Gamma \cup \{\phi\}$ is a set of formulas and if $\mathbf{L}$ is a modal many-valued logic, then $\Gamma \vdash_{\mathbf{L}} \phi$ if and only if there is a finite subset $\{\phi_1, \dots, \phi_r\}$ of Γ and some positive integers $m_1, \dots, m_r$ such that $\vdash_{\mathbf{L}} \phi_1^{m_1} \odot \cdots \odot \phi_r^{m_r}$.*

4. MODAL MANY-VALUED ALGEBRAS AND THE ALGEBRAIC SEMANTIC

We introduce very briefly the varieties of *modal many-valued algebras* and state the completeness result for modal many-valued logics and algebras. This somehow obvious result can be seen as a step towards the construction of the *canonical model* and a possible completeness

theorem for many-valued KRIPKE models. We refer the reader to [5] or [9] for an introduction to the variety of MV-algebras.

Definition 4.1. If $\mathbf{L}$ is a modal many-valued logic then an $\mathbf{L}$ -algebra is an algebra A over the language $\mathcal{L}_{MMV} = \{\rightarrow, \neg, \Box, 0, 1\}$ that satisfies the equations naturally induced by the formulas of $\mathbf{L}$. We denote by $\mathcal{MMV}$ (resp. $\mathcal{MMV}_n$) the variety of $\mathbf{K}$ -algebras (resp. the variety of $\mathbf{K}_n$ -algebras). Members of $\mathcal{MMV}$ (resp. $\mathcal{MMV}_n$) are simply called *modal many-valued algebras* or MMV-algebras (resp. *modal $n+1$ -valued algebras* or $\mathcal{MMV}_n$ -algebras).

A modal many-valued logic $\mathbf{L}$ is often given by a set Γ of axioms, i.e. $\mathbf{L}$ is the smallest modal many-valued logic that contains $\mathbf{K} \cup \Gamma$, and is denoted by $\mathbf{K} + \Gamma$.

Since the most commonly used axiomatization of the variety $\mathcal{MV}$ of MV-algebras is given over the language $\{\oplus, \odot, \neg, 0, 1\}$, we preferably use this language instead of $\mathcal{L}_{MV}$ (with the help of the theorem $(\phi \oplus \psi) \leftrightarrow (\neg\phi \rightarrow \psi)$). Thus, an MMV-algebra is simply an algebra $A = \langle A, \oplus, \odot, \neg, \Box, 0, 1 \rangle$ of type $(2, 2, 1, 1, 0, 0)$ such that

- the reduct of A to the language $\{\oplus, \odot, \neg, 0, 1\}$ is an MV-algebra (i.e. A satisfies the equations $\neg\neg x = x$, $x \oplus 1 = 1$, $\neg 0 = 1$, $x \odot y = \neg(\neg x \oplus \neg y)$, $(x \odot \neg y) \oplus y = (y \odot \neg x) \oplus x$);
- the algebra A satisfies the equations $\Box(x \rightarrow y) \rightarrow \Box x \rightarrow \Box y = 1$, $\Box(x \oplus x) = \Box x \oplus \Box x$, $\Box(x \odot x) = \Box x \odot \Box x$ and $\Box(x \oplus x^m) = (\Box x \oplus (\Box x)^m)$ for every positive integer m .

Similarly, an $\mathcal{MMV}_n$ -algebra is an MMV-algebra whose reduct to the language of MV-algebras is a member of the variety $\mathbf{HSP}(\mathbf{L}_n) = \mathbf{ISP}(\mathbf{L}_n)$.

Recall that on an MV-algebra A , the relation $\leq$ defined by

$$x \leq y \text{ if } x \rightarrow y = 1$$

is a lattice order on A with $x \vee y = (x \rightarrow y) \rightarrow y$ and $x \wedge y = \neg(\neg x \vee \neg y)$.

It is not the object of this paper to study the varieties of MMV-algebras in details. This work should be done in a forthcoming paper. Let us note that the duality developed in [17] for each of the varieties $\mathcal{MMV}_n$ could be used as a tool for the investigation of these varieties.

Definition 4.2. A *filter* of an MMV-algebra A is a filter of its MV-algebra reduct (i.e. a non empty increasing subset of A that contains y whenever it contains x and $x \rightarrow y$). If X is a subset of A , we denote by $\langle X \rangle$ the filter generated by X .

We denote by $\mathcal{F}_{\mathbf{L}}$ the free $\mathbf{L}$ -algebra over the set $\mathbf{Prop}$ of variables for any modal many-valued logic $\mathbf{L}$, i.e. the set of formulas modulo $\mathbf{L}$ -equivalence. In the following, we do not distinguish a formula ϕ with its class $\phi^{\mathbf{L}}$ in $\mathcal{F}_{\mathbf{L}}$.

Recall that the lattice of filters of an MV-algebra A is isomorphic to the lattice of congruences of A . The congruence θ_F associated to a filter F by this isomorphism is defined by $(x, y) \in \theta_F$ if $(x \rightarrow y) \odot (y \rightarrow x) \in F$. As usual, we denote by A/F the quotient A/θ_F .

For our purpose, the next result is fundamental, albeit an obvious one.

Proposition 4.3. *If $\mathbf{L}$ is a modal many-valued logic, and $\Gamma \cup \{\phi\}$ is a set of formulas then $\Gamma \vdash_{\mathbf{L}} \phi$ if and only if $\phi = 1$ in $\mathcal{F}_{\mathbf{L}}/\langle \Gamma \rangle$ or equivalently if ϕ is satisfied in every $\mathbf{L}$ -algebra that satisfies the axioms of Γ .*

5. CONSTRUCTION OF THE CANONICAL MODEL

Here is one of the main contributions of the paper. Recall that the variety of MV-algebras is the variety generated by the algebra $\langle [0, 1], \oplus, \odot, \neg, 0, 1 \rangle$ defined on the real unit interval $[0, 1]$ by $x \oplus y = \min(x + y, 1)$ and $\neg x = 1 - x$ and that an MV-algebra A is simple if and only if it is an isomorphic copy of a subalgebra of $[0, 1]$. Moreover, two isomorphic subalgebras of $[0, 1]$ are necessarily equal (and the isomorphism is the identity). We can thus state the following lemma, which will enable us to define a valuation on the canonical model. A filter of an MV-algebra A is *maximal* if it is maximal among the proper filters of A .

Lemma 5.1. *A filter F of an MV-algebra A is maximal if and only if there is a unique embedding $f : A/F \rightarrow [0, 1]$.*

The idea of the construction of the canonical model for a logic $\mathbf{L}$ is classical. The universe of the model is the set of the maximal filters of $\mathcal{F}_{\mathbf{L}}$ (it coincides with the maximal consistent extensions of $\mathbf{L}$). But, in order to simplify the expression of our results, it is better to identify, with the help of Lemma 5.1, the set of the maximal filters of $\mathcal{F}_{\mathbf{L}}$ with the set $\mathcal{MV}(\mathcal{F}_{\mathbf{L}}, [0, 1])$ of the homomorphisms of MV-algebras from $\mathcal{F}_{\mathbf{L}}$ to $[0, 1]$.

Definition 5.2. The *canonical model* for a modal many-valued logic $\mathbf{L}$ is the model $\mathcal{M}_{\mathbf{L}} = (W_{\mathbf{L}}, R_{\mathbf{L}}, \text{Val}_{\mathbf{L}})$ where

- the universe $W_{\mathbf{L}}$ of $\mathcal{M}_{\mathbf{L}}$ is the set $\mathcal{MV}(\mathcal{F}_{\mathbf{L}}, [0, 1])$;
- the binary relation $R_{\mathbf{L}}$ is defined by

$$(u, v) \in R_{\mathbf{L}} \text{ if } \forall \phi \in \mathcal{F}_{\mathbf{L}} (u(\Box\phi) = 1 \Rightarrow v(\phi) = 1),$$

- the valuation map $\text{Val}_{\mathbf{L}} : \text{Prop} \times W_{\mathbf{L}} \rightarrow [0, 1]$ is defined by

$$\text{Val}_{\mathbf{L}}(u, p) = u(p).$$

Note that the definition of the canonical model for $\mathbf{K}_1$ coincides with the classical definition of the canonical model for the Boolean basic modal logic (if we identify any maximal filter F of $\mathcal{F}_{\mathbf{K}_1}$ with the quotient map $\pi_F : \mathcal{F}_{\mathbf{K}_1} \rightarrow \{0, 1\}$ and if we identify the Boolean valuation map $\text{Val} : \text{Prop} \rightarrow \mathcal{P}(W_{\mathbf{K}_1})$ with its characteristic function).

The main result of this section is that the map $\text{Val}_{\mathbf{L}}$ extends to formulas. Before considering the proof of this result, we need the following definition.

Definition 5.3. We denote by $\mathbb{D}$ the subset of $\mathbb{Q}$ that contains the numbers that can be written as a finite sum of powers of 2. If r is in $\mathbb{D} \cap [0, 1]$, we denote by τ_r a composition of the terms $x \oplus x$ and $x \odot x$ such that $\tau_r(x) < 1$ for every $x \in [0, r[$ and $\tau_r(x) = 1$ for every $x \in [r, 1]$. A proof of the existence of such terms can be found in [16] for example. Furthermore, we can always choose τ_r such that $\tau_r(x) = 1$ for every $x \in \mathbb{L}_n \cap [r, 1]$ (but this choice is not independant of n).

Lemma 5.4. *If $\mathbf{L}$ is a modal many-valued logic and if $u, v \in W_{\mathbf{L}}$, then $(u, v) \in R_{\mathbf{L}}$ if and only if $u \circ \Box \leq v$.*

Proof. The right to left part of the assertion is clear. Let us prove the left to right part and suppose that there is a ϕ in $\mathcal{F}_{\mathbf{L}}$, a v in $R_{\mathbf{L}}u$ and an r in $\mathbb{D} \cap [0, 1]$ such that $v(\phi) < r \leq u(\Box\phi)$.

It follows that

$$\tau_r(v(\phi)) = v(\tau_r(\phi)) < 1 \quad \text{and} \quad 1 = \tau_r(u(\Box\phi)) = u(\tau_r(\Box\phi)) = u(\Box\tau_r(\phi)),$$

which is a contradiction since $uR_{\mathbf{L}}v$. □

Proposition 5.5. *If $\mathbf{L}$ is a modal many-valued logic, then*

$$\text{Val}_{\mathcal{M}_{\mathbf{L}}}(\phi, u) = u(\phi)$$

for any ϕ in $\mathcal{F}_{\mathbf{L}}$ and u in $W_{\mathbf{L}}$.

Proof. The non trivial statement is the equality

$$(5.1) \quad u(\Box\phi) = \bigwedge_{v \in R_{\mathbf{L}}u} v(\phi).$$

The inequality $\leq$ is the content of Lemma 5.4.

Suppose now that the equality does not hold in (5.1), but just the strict inequality $<$. Then, there is an r in $\mathbb{D} \cap [0, 1]$ such that

$$u(\Box\phi) < r \leq \bigwedge_{v \in R_{\mathbf{L}}u} v(\phi).$$

It means that for any $v \in R_{\mathbf{L}}u$, the maximal filter $v^{-1}(1)$ of $\mathcal{F}_{\mathbf{L}}$ contains $\tau_r(\phi)$ and that the filter $\Box^{-1}u^{-1}(1)$ does not contain $\tau_r(\phi)$.

But, since $(u, v) \in R_{\mathbf{L}}$ if and only if $\Box^{-1}u^{-1}(1) \subseteq v^{-1}(1)$, it follows that the maximal filters of $\mathcal{F}_{\mathbf{L}}$ that contain $\Box^{-1}u^{-1}(1)$ are exactly the $v^{-1}(1)$ with $v \in R_{\mathbf{L}}u$, while each of these maximal filters contains $\tau_r(\phi)$. It means that the class of $\tau_r(\phi)$ in $\mathcal{F}_{\mathbf{L}}/\Box^{-1}u^{-1}(1)$ is infinitely great, so that $\tau_r(\phi) \oplus \tau_r(\phi)^m$ belongs to $\Box^{-1}u^{-1}(1)$ for every positive integer m . It follows that

$$1 = u(\Box(\tau_r(\phi) \oplus \tau_r(\phi)^m)) \leq u(\tau_r(\Box\phi) \oplus (\tau_r(\Box\phi))^m),$$

for any positive integer m , so that $u(\tau_r(\Box\phi))$ is infinitely great in $u(\mathcal{F}_{\mathbf{L}})$. Since $u(\mathcal{F}_{\mathbf{L}})$ is a subalgebra of $[0, 1]$, we obtain that $u(\tau_r(\Box\phi)) = 1$, a contradiction. □

6. COMPLETENESS RESULTS

Proposition 5.5 enables to prove completeness results for some modal many-valued logics... but only for some of them. Indeed, with the help of Proposition 5.5, we obtain that if Γ is a set of axioms then a formula ϕ that is valid in every model of Γ is deducible from Γ if the algebra $\mathcal{F}_{\mathbf{K}}/\langle\Gamma\rangle$ is semi-simple, i.e. if $\langle\Gamma\rangle$ is the only element of $\mathcal{F}_{\mathbf{L}}/\langle\Gamma\rangle$ that is in every maximal filter of $\mathcal{F}_{\mathbf{L}}/\langle\Gamma\rangle$. Unfortunately, contrary to the two-valued case (where every logic is the intersection of its maximal consistent extensions), we can not ensure *a priori* that $\mathcal{F}_{\mathbf{L}}/\langle\Gamma\rangle$ is a semi-simple algebra.

Hopefully, there are some very interesting logics for which the completeness result can be stated.

6.1. Modal finitely-valued logics. The first family of systems that admit the many-valued KRIPKE models as a complete semantic is the finitely-valued ones.

Theorem 6.1. *If $\Gamma \cup \{\phi\}$ is a set of formulas, then $\Gamma \vdash_{\mathbf{K}_n} \phi$ if and only if ϕ is valid in every $n + 1$ -valued KRIPKE model of Γ .*

Proof. The algebra $\mathcal{F}_{\mathbf{K}_n}/\langle \Gamma \rangle$ is a member of $\mathbf{HSP}(\mathbb{L}_n) = \mathbf{ISP}(\mathbb{L}_n)$ and so is semi-simple. $\square$

Note that by considering $n = 1$, the preceding proposition boils down to the completeness result for Boolean basic modal logic and KRIPKE semantic.

We have announced in section 4 the following result which is an application of the preceding completeness theorem.

Proposition 6.2. *If n is a positive integer then $\vdash_{\mathbf{K}_n} \Box(p \wedge q) \leftrightarrow (\Box p \wedge \Box q)$.*

Moreover, we can simplify the axiomatization 3.1 of $\mathbf{K}_n$. We can indeed get rid off the family of axioms that expresses the conservative law of $\Box$ with respect to the infinitely great elements.

Proposition 6.3. *If $\mathbf{MV}_n$ denotes the $n + 1$ -valued ŁUKASIEWICZ logic and if $\mathbf{K}'_n = \mathbf{MV}_n + \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q) + \Box(p \oplus p) \leftrightarrow (\Box p \oplus \Box p) + \Box(p \odot p) \leftrightarrow (\Box p \odot \Box p)$, then $\mathbf{K}_n = \mathbf{K}'_n$.*

Proof. In the proof of Proposition 5.5 with $\mathbf{L} = \mathbf{K}_n$, we can deduce directly that $u(\Box \tau_r(\phi)) = 1$ from the fact that $\tau_r(\phi)$ is infinitely great in $\mathcal{F}_{\mathbf{K}_n}/\Box^{-1}u^{-1}(1)$, since $\mathcal{F}_{\mathbf{K}_n}/\Box^{-1}u^{-1}(1)$ has no non trivial infinitely great element. It means that Proposition 5.5 stands with $\mathbf{L} = \mathbf{K}'_n$ and that $\vdash_{\mathbf{K}'_n} \phi$ for any formula ϕ that is valid in any $n + 1$ -valued KRIPKE model. We can thus conclude since for any positive integer m , the formula $\Box(p \oplus p^m) \rightarrow (\Box p \oplus (\Box p)^m)$ is a tautology. $\square$

Apart from the completeness result, the extensions of $\mathbf{K}_n$ seem to share interesting properties with the Boolean modal logics. For instance, the paper [17] is dedicated to the construction of a duality for $\mathbf{K}_n$ -algebras and a class of topological structures. For $n = 1$, this duality coincides with the STONE duality for modal algebras. The role played by the duality for $\mathbf{K}_n$ -algebras is as important as the role played by the STONE duality for modal algebras in Boolean modal logic. Indeed, the class of the dual structures (called $\mathcal{MX}_n$ -structures) forms a very adequate semantic since any extension $\mathbf{L}$ of $\mathbf{K}_n$ is complete with respect to the class of $\mathcal{MX}_n$ -structures in which $\mathbf{L}$ is valid.

Moreover, the construction of this duality suggests two ways of going from $n + 1$ -valued KRIPKE models to frames (and conversely). Indeed, we can obviously define a frame $\mathfrak{F} = \langle W, R \rangle$ as a set W with a binary relation R on W . Then, a frame $\mathfrak{F} = \langle W, R \rangle$ becomes a model by the addition of a valuation $\text{Val} : \text{Prop} \times W \rightarrow \mathbb{L}_n$. So, the set of truth values in a world w is given by the valuation, at the model level and not at the frame level. Now, it is also possible (and as we shall see, relevant) to consider some new (first order) structures, called $n + 1$ -frames, in which the set of truth values in a world w is known *a priori*, without any reference to a valuation. In the following definition, we denote by $\text{div}(n)$ the set of the positive divisors of n .

Definition 6.4. An $n + 1$ -frame $\mathfrak{F} = \langle W, \{r_m \mid m \in \text{div}(n)\}, R \rangle$ is given by a set W , a subset r_m of W for every m in $\text{div}(n)$ and a relation $R \subseteq W \times W$ such that

- (1) for every m and k in $\text{div}(n)$, the intersection $r_m \cap r_k$ coincides with $r_{\text{gcd}(m,k)}$ and $r_n = W$;
- (2) for every m in $\text{div}(n)$, the set $Rr_m = \{w \mid \exists w' \in r_m w'Rw\}$ of the successors of the elements of r_m is a subset of r_m .

A model $\langle W', R', \text{Val} \rangle$ is *based on an the $n+1$ -frame* $\mathfrak{F} = \langle W, \{r_m \mid m \in \text{div}(n)\}, R \rangle$ if $W = W'$, $R = R'$ and $\text{Val}(p, w) \in \mathbb{L}_m$ for any m in $\text{div}(n)$, any w in r_m and p in Prop .

Validity in $n+1$ -frames is defined similarly as in the class of frames.

Thus, an $n+1$ -frame is obtained from a frame by restricting the class of valuations that can be added to this frame to define an $n+1$ -KRIPKE model. We should so have a gain in the expressivity of the class of $n+1$ -frames with regard to the class of frames. For instance, there are some extensions $\mathbf{L}$ of $\mathbf{K}_n$ that are characterized by a class of $n+1$ -frames but that are not characterized by any class of frames. Here are a few easy examples.

Definition 6.5. A modal many-valued logic $\mathbf{L}$ is *KRIPKE complete* (resp. *tabular*) if there is a class of frames K (resp. a finite frame $\mathfrak{F}$) such that $\mathbf{L}$ is the set of formulas that are valid in every frame of K (resp. in $\mathfrak{F}$).

Similarly, a logic $\mathbf{L}$ is *$n+1$ -KRIPKE complete* (resp. *$n+1$ -tabular*) if there is a class of $n+1$ -frames (resp. an $n+1$ -frame $\mathfrak{F}$) such that $\mathbf{L}$ is the set of formulas that are valid in every model based on a frame of K (resp. on $\mathfrak{F}$).

If $\mathfrak{F} = \langle W, \{r_m \mid m \in \text{div}(n)\}, R \rangle$ and $\mathfrak{F}' = \langle W', \{r'_m \mid m \in \text{div}(n)\}, R' \rangle$ are two $n+1$ -frames, a map $f : W \rightarrow W'$ is called a *$n+1$ - π -morphism* if the three following conditions are satisfied:

- (1) if u and v are in W and $(u, v) \in R$ then $(f(u), f(v)) \in R'$;
- (2) if $u \in W$ and $v' \in W'$ with $(f(u), v') \in R$ then there exists a v in Ru such that $f(v) = v'$;
- (3) if $u \in r_m$ then $f(u) \in r'_m$.

We leave to the reader the task to prove that validity is preserved under $n+1$ - π -morphic image, i.e. that if $f : \mathfrak{F} \rightarrow \mathfrak{F}'$ is a surjective $n+1$ - π -morphism between two $n+1$ -frames $\mathfrak{F}$ and $\mathfrak{F}'$, then $\mathfrak{F}' \models \phi$ whenever $\mathfrak{F} \models \phi$.

Proposition 6.6. *Assume that $n \geq 2$. We have the following completeness results.*

- (1) *The logic $\mathbf{L}_1 = \mathbf{K}_n + (\Box p \vee \Box \neg p)$ is $n+1$ -KRIPKE complete with respect to the class of the $n+1$ -frames that satisfy $\forall u Ru \subseteq r_1$ but is not KRIPKE-complete.*
- (2) *The logic $\mathbf{L}_2 = \mathbf{K}_n + \Box(p \vee \neg p) + \Box \Box p + \neg(\Diamond p \wedge \Diamond \neg p)$ is $n+1$ -tabular but is not even KRIPKE complete.*
- (3) *The logic $\mathbf{K}_n + \Box x \rightarrow x$ is KRIPKE-complete with respect to the class of reflexive frames.*
- (4) *The logic $\mathbf{K}_n + \Box x \rightarrow \Box \Box x$ is KRIPKE-complete with respect to the class of transitive frames.*

Proof. (1) It is clear, by the definition of a model based on an $n+1$ -frame that $\mathbf{L}_1$ is characterized by the class of $n+1$ -frames that satisfy the first order formula $\forall u Ru \subseteq r_1$.

Suppose then that K is a class of frames that characterizes $\mathbf{L}_1$. First note that we may suppose that K contains a frame $\langle W, R \rangle$ with R non trivial. Otherwise, for any frame $\mathfrak{F}$ of K

and any formula ϕ , the formula $\Box\phi$ is valid in $\mathfrak{F}$ and is so a theorem of $\mathbf{L}_1$. But it is easy to construct a $\mathbf{L}_1$ -counter-model for $\Box(p \wedge \neg p)$.

Now, if $\mathfrak{F} = \langle W, R \rangle$ is a frame of K with a non trivial relation R and if $\mathcal{M} = \langle W, R, \text{Val} \rangle$ is a model based on $\mathfrak{F}$ and $w, v \in W$ with wRv , it follows that $\mathcal{M}, w \models (\Box p \vee \Box \neg p)$. We deduce that $\text{Val}(p, v) \in \{0, 1\}$. Then, if we denote by $\mathcal{M}' = \langle W, R, \text{Val}' \rangle$ the model based on $\mathfrak{F}$ defined by

$$\text{Val}'(q, u) = \begin{cases} \text{Val}(q, u) & \text{if } q \neq p \text{ or } u \neq v, \\ \frac{1}{n} & \text{if } q = p \text{ and } u = v, \end{cases}$$

it appears that $\Box p \vee \Box \neg p$ is not true in $\mathcal{M}'$, a contradiction since $\mathcal{M}'$ is based on a frame of K .

(2) Let us consider the $n + 1$ -frame $\mathfrak{F}$ whose universe is $\{u, v\}$ with $(u, v) \in R$, $u \in r_n$ and $v \in r_1$ (we only specify for any world of an $n + 1$ -frame the smallest of the subsets r_m that contain this world) and the $n + 1$ -frame $\mathfrak{F}'$ as the one irreflexive point belonging to r_n .

It is clear that any formula of $\mathbf{L}_2$ is satisfied in $\mathfrak{F}$ and $\mathfrak{F}'$. Now, suppose that ϕ is a formula that is satisfied in $\mathfrak{F}$ and $\mathfrak{F}'$ and prove that ϕ belongs to $\mathbf{L}_2$. It suffices to prove that ϕ is valid in the canonical model of $\mathbf{L}_2$.

First, if w is a world of $\mathcal{M}_{\mathbf{L}_2}$, then $w \models \Box\Box\psi$ for every formula ψ and thus the subframe $R_{\mathbf{L}_2}^\omega w$ of $\langle W_{\mathbf{L}_2}, R_{\mathbf{L}_2} \rangle$ generated by w is equal to $R_{\mathbf{L}_2} w$ and $(w, w) \notin R_{\mathbf{L}_2}$.

Then, since $w \models \neg(\Diamond\phi \wedge \Diamond\neg\phi)$ for every formula ϕ , it follows that $|R_{\mathbf{L}_2}| \leq 1$. Otherwise there are two worlds t and s in $R_{\mathbf{L}_2} w$. Since we work in the canonical model, it means that there is a formula ψ such that $t(\psi) \neq s(\psi)$. We deduce that $w \not\models \neg(\Diamond\psi \wedge \Diamond\neg\psi)$.

Now, with the help of the axiom $\Box(p \vee \neg p)$, we obtain that $R_{\mathbf{L}_2}^\omega w$ is an $n + 1$ - π -morphic image of $\mathfrak{F}$ or $\mathfrak{F}'$ and thus $\mathcal{M}_{\mathbf{L}_2}, w \models \phi$.

We leave to the reader the task to prove, similarly as in (1), that the logic $\mathbf{L}_2$ is not KRIPKE complete.

For (3) and (4), it suffices to mimic the classical proofs. □

The preceding proposition illustrates the difference between frame definability and $n + 1$ -frame definability and gives a first simple example of correspondence theory for modal many-valued formulas and $n + 1$ -frames. We should study in more details in a forthcoming paper this dissimilarity in frame definability.

6.2. Infinitary modal many-valued systems. Another way of obtaining completeness results is to extend the modal systems that we have defined by an infinitary deduction rule.

Definition 6.7. The *infinitary modal many-valued system* has the set of well-formed formulas, the *modus ponens*, the necessitation rule and the rule

$$(\text{Inf}) \quad \frac{\phi \oplus \phi, \phi \oplus \phi^2, \dots, \phi \oplus \phi^n, \dots}{\phi}$$

as deduction rules.

If Γ is a set of axioms and ϕ is a formula, we write $\Gamma \vdash_\infty \phi$ if ϕ appears in a possibly infinite sequence $(\psi_\beta)_{\beta \leq \alpha}$ of formulas that belongs to Γ , are obtained by substitution in a formula belonging to Γ or that are obtained by the application of a deduction rule from previous formulas of the sequence.

Note that if ϕ is a formula and $\Gamma \vdash_{\mathbf{K}} \phi$ then $\Gamma \vdash_{\infty} \phi$. We can then state the following completeness result.

Proposition 6.8. *If $\Gamma \cup \{\phi\}$ is a set of formulas, then $\Gamma \vdash_{\infty} \phi$ if and only if ϕ is true in every many-valued KRIPKE model of Γ .*

Proof. If $\Gamma \vdash_{\infty} \phi$, the result follows from that fact that there is no infinitely great element in the MV-algebra $[0, 1]$, so that the rule (Inf) preserves tautologies.

Now, if ϕ is true in any model of Γ , then ϕ is infinitely great in the algebra $\mathcal{F}_{\mathbf{K}}/\langle\Gamma\rangle$, which means that for any integer $m \geq 2$, the element $\phi \oplus \phi^m$ is equal to $\langle\Gamma\rangle$ in $\mathcal{F}_{\mathbf{K}}/\langle\Gamma\rangle$, or equivalently that $\Gamma \vdash_{\mathbf{K}} \phi \oplus \phi^m$ for any integer $m \geq 2$. We can conclude using the rule (Inf). $\square$

7. CONCLUSIONS

We propose some tracks than could be followed in the future.

- Infinitary vs finitary modal systems. The general completeness result that is proposed in this paper involves an infinitary deduction rule. Since we can get rid off this rule in the case of the finitely-valued modal logics, the question to determine the minimal extensions of $\mathbf{K}$ for which the completeness result – without the infinitary rule – can be stated should be considered. We do not know if $\vdash_{\infty} \phi$ is equivalent to $\vdash_{\mathbf{K}} \phi$.
- Varieties of MMV-algebras and MMV $_n$ -algebras. We have not given any significant information about these varieties. A good tool for the studies of varieties of MMV $_n$ -algebras could be the topological duality constructed in [17]. A problem that could be solved in this way is the characterization of finitely generated algebras (following some ideas of [6]).
- KRIPKE-completeness, $n + 1$ -KRIPKE-completeness and correspondence theory: we should study in detail, with the tool of universal algebra and canonical extensions, the dissimilarity between KRIPKE-completeness and $n + 1$ -KRIPKE-completeness and consider the problem of the correspondence between modal many-valued formulas and first order sentences on frames and $n + 1$ -frames.
- Temporal logic and propositional dynamic logic: we should give interest to the construction of some particular systems of modal $n + 1$ -valued logics such as $n + 1$ -valued temporal logic or dynamic logic. By their nature, such systems could be useful for computer scientists for example.
- Extension to more general languages. The results of this paper are obtained for the basic modal language with one unary modal operator. They should be extended to languages containing k -ary modalities.

REFERENCES

- [1] P. Blackburn, M. de Rijke, and Y. Venema. *Modal logic*, volume 53 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, Cambridge, 2001.
- [2] A. Chagrov and M. Zakharyashev. *Modal logic*, volume 35 of *Oxford Logic Guides*. The Clarendon Press Oxford University Press, New York, 1997. Oxford Science Publications.
- [3] C. C. Chang. Algebraic analysis of many valued logics. *Trans. Amer. Math. Soc.*, 88:467–490, 1958.
- [4] C. C. Chang. A new proof of the completeness of the Łukasiewicz axioms. *Trans. Amer. Math. Soc.*, 93:74–80, 1959.

- [5] R. L. O. Cignoli, I. M. L. D'Ottaviano, and D. Mundici. *Algebraic foundations of many-valued reasoning*, volume 7 of *Trends in Logic—Studia Logica Library*. Kluwer Academic Publishers, Dordrecht, 2000.
- [6] L. Esakia and R. Grigolia. The criterion of Brouwerian and closure algebras to be finitely generated. *Polish Acad. Sci. Inst. Philos. Sociol. Bull. Sect. Logic*, 6(2):46–52, 1977.
- [7] M. Fitting. Many-valued modal logics. II. *Fund. Inform.*, 17(1-2):55–73, 1992.
- [8] M. C. Fitting. Many-valued modal logics. *Fund. Inform.*, 15(3-4):235–254, 1991.
- [9] J. Gispert and D. Mundici. MV-algebras: a variety for magnitudes with Archimedean units. *Algebra Universalis*, 53(1):7–43, 2005.
- [10] S. Gottwald. *A treatise on many-valued logics*, volume 9 of *Studies in Logic and Computation*. Research Studies Press Ltd., Baldock, 2001.
- [11] B. Jónsson and A. Tarski. Boolean algebras with operators. I. *Amer. J. Math.*, 73:891–939, 1951.
- [12] B. Jónsson and A. Tarski. Boolean algebras with operators. II. *Amer. J. Math.*, 74:127–162, 1952.
- [13] S. A. Kripke. Semantical analysis of modal logic. I. Normal modal propositional calculi. *Z. Math. Logik Grundlagen Math.*, 9:67–96, 1963.
- [14] J. Łukasiewicz. O logice trójwartościowej. *Ruch Filozoficzny*, 5:170–171, 1920.
- [15] J. Łukasiewicz. *Selected works*. North-Holland Publishing Co., Amsterdam, 1970. Edited by L. Borkowski, Studies in Logic and the Foundations of Mathematics.
- [16] P. Ostermann. Many-valued modal propositional calculi. *Z. Math. Logik Grundlagen Math.*, 34(4):343–354, 1988.
- [17] B. Teheux. A duality for the algebras of a Łukasiewicz $n + 1$ -valued modal system. *To appear in Studia Logica*, 2006.

E-mail address: `g.hansoul@ulg.ac.be`, `b.teheux@ulg.ac.be`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE LIÈGE, 12, GRANDE TRAVERSE, 4000 LIÈGE, BELGIUM.